

A DIRECT METHOD FOR OBTAINING THE DIFFERENTIAL CONSERVATION LAWS

A.A. CHERNITSKII

*St. Petersburg Electrotechnical University, Prof. Popov str. 5,
St. Petersburg 197376, Russia ; E-mail: aa@cher.etu.spb.ru*

Abstract

A direct method for obtaining the differential conservation laws in the field theory from the principle of stationary action is proposed. The method is based on a variation of field functions through small local transformation of a special kind. The action of general theory of relativity are considered.

According to the known Noether's theorem ¹ the existence of the differential conservation laws in the field theory is connected with invariance of the action under continuous transformations of dependent and independent variables. If we have such transformation, then the Noether's theorem gives a form of appropriate differential conservation law. Another methods for obtaining differential conservation laws also are used. For example, it is of common knowledge that the conservation law of the impulse-energy tensor can be easily obtained from the Euler's system of equations ². A comparatively simple method of obtaining the conservation laws from the principle of stationary action will be proposed here. We will derive necessary conditions of the action stationarity (variation of the action is zero) that can take the form of differential conservation laws.

Let us consider the following action in n -dimensional space.

$$S = \int_{\Omega} \mathcal{L}(A_{\mu}, B_{\nu\mu}, x^{\nu}) (dx)^n \quad (1)$$

where $A_{\mu} = A_{\mu}(\{x^{\nu}\})$ are tensor components of the field,
 $B_{\nu\mu}$ is a derivative of the function $A_{\mu}(\{x^{\nu}\})$ by its ν -th argument,
 Ω is a region of space, $(dx)^n \equiv dx^1 dx^2 \dots dx^n$.

As it is well known, direct variation of the functions $A_{\mu}(x)$ generate a Euler's system of equations as condition of the stationary action (1). However, it is possible to obtain a variation of the functions $A_{\mu}(x)$ through a small local transformation of some special kind. For example we can use a small local shift of whole field configuration. Let the functions $A_{\mu}(\{x^{\nu}\})$ provide the stationarity of the action S (1) ($\delta S = 0$). Let us consider varied functions $A_{\mu}(\{x^{\nu} - \varepsilon \delta a^{\nu}(x)\})$. Here $\delta a^{\nu}(x) = 0$ on the border of region Ω . So we have the variation of the field functions with help of a small shift of the field configuration in the space, the shift size being function of the coordinates. Thus the coordinate system does not vary, it is only the functional dependence that varies. In any fixed point of region Ω this shift generates the variation of values of the functions $A_{\mu}(x)$ in general case. The functions can take any values between minimum and maximum of the stationary configuration of the field ($\delta S = 0$). Thus, this is variation of the functions $A_{\mu}(x)$

but with a specific way of variation. It is evident that such variation restricts the set of the comparison functions in general case. Thus we can derive only necessary conditions of the stationary action with help of this variation; that is, if $\delta S = 0$, then we have certain conditions for the dependent variables $A_\mu(x)$.

We shall substitute functions $A_\mu(\{x^\nu - \varepsilon \delta a^\nu(x)\})$ into Lagrangian. Now it is necessary to extract the linear terms with δa^ν . In the regular way, we differentiate the action by ε and we take $\varepsilon = 0$ for this purpose. Thus we have the variation of the action in the following form.

$$\delta S = \int_{\Omega} \left(f^{\mu\nu} B_{\rho\nu} \frac{\partial \delta a^\rho}{\partial x^\mu} + \frac{\partial \mathcal{L}}{\partial A_\nu} B_{\rho\nu} \delta a^\rho + f^{\mu\nu} \frac{\partial B_{\mu\nu}}{\partial x^\rho} \delta a^\rho \right) (dx)^n \quad (2)$$

With the partial integration and taking into account that $\delta a^\rho(x)$ are arbitrary functions, we have the necessary condition of the stationary action in the following form.

$$\frac{\partial}{\partial x^\mu} (f^{\mu\nu} B_{\rho\nu}) - \frac{\partial \mathcal{L}}{\partial A_\nu} B_{\rho\nu} - f^{\mu\nu} \frac{\partial B_{\mu\nu}}{\partial x^\rho} = 0 \quad (3)$$

If Lagrangian does not depend explicitly on the coordinates x^ν , the relation (3) can be written in the following conservation law form.

$$\frac{\partial}{\partial x^\mu} (f^{\mu\nu} B_{\rho\nu} - \mathcal{L} \delta_\rho^\mu) = 0 \quad (4)$$

For space-time case this is the known conservation law of energy-impulse tensor.

In the same way we can obtain the variations of the functions $A_\mu(\{x^\nu\})$ through small local turn about a coordinate center. For simplicity let us consider $A_\mu(\{x^\nu\})$ as components of vector field. Then, if $A_\mu(\{x^\nu\})$ provide the stationary action, the varied functions will have the form $L_{\cdot\mu}^\rho A_\rho(\{L_{\cdot\delta}^\nu x^\delta\})$, where $L_{\cdot\mu}^\rho = \delta_\mu^\rho + \varepsilon \delta L_{\cdot\mu}^\rho(x)$. $\delta L^{\rho\mu}(x) = -\delta L^{\mu\rho}(x)$ are turn parameters. Also we assume that $\delta L_{\cdot\mu}^\rho(x) = 0$ on the border. Here the variation of the arguments of the field functions generates a shift, where the shift size is proportional to the distance from the coordinate center. Substituting the varied functions into Lagrangian, differentiating action by ε and taking $\varepsilon = 0$ we obtain variation of the action in the following form.

$$\delta S = \int_{\Omega} \left(f^{\mu\nu} B_{\rho\nu} x^\beta \frac{\partial \delta L_{\cdot\beta}^\rho}{\partial x^\mu} + f^{\mu\nu} A_\rho \frac{\partial \delta L_{\cdot\nu}^\rho}{\partial x^\mu} + \frac{\partial \mathcal{L}}{\partial A_\nu} B_{\rho\nu} x^\beta \delta L_{\cdot\beta}^\rho + \right. \quad (5) \\ \left. + f^{\mu\nu} \frac{\partial B_{\mu\nu}}{\partial x^\rho} x^\beta \delta L_{\cdot\beta}^\rho \right) (dx)^n$$

When we obtained (5) we took into account that Lagrangian is invariant under turn transformations with constant parameters. We use partial integration to (5). Using the antisymmetry of $\delta L^{\rho\mu}$ and the arbitrariness of the turn parameters we obtain the necessary condition of the stationary action in the following form.

$$\frac{\partial}{\partial x^\mu} \left[f^{\mu\nu} (B_{\rho\nu} x_\beta - B_{\beta\nu} x_\rho) + (f_{\cdot\beta}^\mu A_\rho - f_{\cdot\rho}^\mu A_\beta) \right] - \quad (6) \\ - \frac{\partial \mathcal{L}}{\partial A_\nu} (B_{\rho\nu} x_\beta - B_{\beta\nu} x_\rho) - f^{\mu\nu} \left(\frac{\partial B_{\mu\nu}}{\partial x^\rho} x_\beta - \frac{\partial B_{\mu\nu}}{\partial x^\beta} x_\rho \right) = 0$$

If Lagrangian does not depend explicitly on the coordinates x^ν , relation (6) can be written in the following conservation law form.

$$\frac{\partial}{\partial x^\mu} \left[f^{\mu\nu} (B_{\rho\nu} x_\beta - B_{\beta\nu} x_\rho) + (f_{\beta}^\mu A_\rho - f_{\rho}^\mu A_\beta) - \mathcal{L} (\delta_\rho^\mu x_\beta - \delta_\beta^\mu x_\rho) \right] = 0 \quad (7)$$

For space-time case this is the known conservation law of angular momentum tensor.

We shall consider now the variation of field functions through a local transformation of more general kind than the turn. We consider the known Lagrangian that gives the Einstein's equations for free gravitational field ³.

$$\mathcal{L} = g^{\mu\nu} (\Gamma_{\mu\sigma}^\rho \Gamma_{\nu\rho}^\sigma - \Gamma_{\mu\nu}^\rho \Gamma_{\rho\sigma}^\sigma) \sqrt{|g|} \quad ; \quad \Gamma_{\mu\nu}^\rho \equiv \frac{1}{2} g^{\rho\gamma} (B_{\nu\gamma\mu} + B_{\mu\gamma\nu} - B_{\gamma\mu\nu}) \quad (8)$$

where $B_{\nu\gamma\mu}$ is a derivative of function $g_{\gamma\mu}(\{x^\nu\})$ with its ν -th argument.

The action with this Lagrangian is invariant for the transformation $x^\rho \rightarrow G_{\cdot\mu}^\rho x^\mu$, where all components of the matrix $G_{\cdot\mu}^\rho$ are independent and constant. Let us consider a variation of the metric tensor components through the transformation $g_{\mu\delta}(\{x^\nu\}) \rightarrow G_{\cdot\mu}^\rho G_{\cdot\delta}^\gamma g_{\rho\gamma}(\{G_{\cdot\sigma}^\nu x^\sigma\})$, where $G_{\cdot\mu}^\rho = G_{\cdot\mu}^\rho(x)$. Then a variation of the action has the following form.

$$\begin{aligned} \delta S = & \int_{\Omega} \left(\mathcal{L} \delta G_{\cdot\rho}^\rho + f^{\mu\nu\sigma} B_{\rho\nu\sigma} x^\beta \frac{\partial \delta G_{\cdot\beta}^\rho}{\partial x^\mu} + 2 f^{\mu\nu\sigma} g_{\rho\sigma} \frac{\partial \delta G_{\cdot\nu}^\rho}{\partial x^\mu} + \right. \\ & \left. + \frac{\partial \mathcal{L}}{\partial g_{\mu\nu}} B_{\rho\mu\nu} x^\beta \delta G_{\cdot\beta}^\rho + f^{\mu\nu\sigma} \frac{\partial B_{\mu\nu\sigma}}{\partial x^\rho} x^\beta \delta G_{\cdot\beta}^\rho \right) (dx)^n \quad ; \quad f^{\mu\nu\sigma} \equiv \frac{\partial \mathcal{L}}{\partial B_{\mu\nu\sigma}} \end{aligned} \quad (9)$$

As we see, in this case the necessary condition of the stationary action can be written in the following conservation law form.

$$\frac{\partial}{\partial x^\mu} (f^{\mu\nu\sigma} B_{\rho\nu\sigma} x^\beta + 2 f^{\mu\beta\sigma} g_{\rho\sigma} - \mathcal{L} x^\beta \delta_\rho^\mu) = 0 \quad (10)$$

So we consider some transformation of dependent and independent variables with continuous parameters. Then we derive the necessary conditions of the stationary action, the variation of the field functions being obtained through this transformation but we take its parameters as depending on the independent variables. Thus we have the necessary conditions of the stationary action of a special kind. In certain cases this conditions can be written in a differential conservation laws form. These cases are specified by Noether's theorem but the presented direct method more conveniently sometimes to use.

References

1. R. Curant and D. Hilbert, Method der Mathematical Physik (Verlag von Julius Springer, Berlin, 1931).
2. L. Landau and E. Lifshitz, The Classical Theory of Fields (Pergamon, Elmsford, 4th ed., 1975).
3. P.A.M. Dirac, General Theory of Relativity (John Wiley & Sons, New York, 1975).